CASE AND CONTACT MANAGEMENT

2022-03-23

	STAFF IN EDUCATION/HIGH SCHOOL STUDENTS/GENERAL POPULATION OVER 12 YRS		
	PROTECTED Worker is considered protected: If having had COVID-19 after December 20 th 2022	VACCINATED* OR WITH HISTORY OF COVID-19 prior to December 20 th *at least one dose	UNVACCINATED WITH NO HISTORY OF COVID-19
1. POSITIVE	N/A – SEE BELOW 'COVID SYMPTOMS'	 5-day isolation If symptoms decreasing and no fever x 24 hrs, can return to activities Must protect others by: Wearing mask at all times, Staying away from vulnerable people, Staying 2 meters from others at all times 	 10-day isolation period 21-day isolation for immunosuppressed people, or anyone who was hospitalized (ICU) for COVID-19. *10-day isolation can be reduced if: 5 full days have passed AND On day 6 symptoms have diminished and no fever X24 hours without medication AND With a negative rapid test result
2. COVID SYMPTOMS	 Withdraw from work until symptoms resolve Consider other diagnosis/ test for that diagnosis Apply IP&C* measures for the alternate diagnosis No Rapid or PCR test needed (could test + for up to 6 months after a previous diagnosis of COVID-19) 	 Isolate immediately Perform a rapid test: If negative: continue to isolate and perform a second test 24-36 hours later, if still negative and symptoms resolved without fever, can return to normal activities, If positive: continue to isolate and follow guidelines in section 1 If able to access a PCR test: If negative: return to normal activities, If positive: continue to isolate and follow guidelines in section 1 	
3. ASYMPTOMATIC CONTACT OF A POSITIVE CASE	 No isolation – Can work No screening or PCR required Consider other diagnosis/ test for that diagnosis Apply IP&C measures for the alternate diagnosis 	FOR ALL TYPES OF CONTACT FOR 10 DAYS AFTER THE LAST CONTACT WITH A POSITIVE CASE: Can go to work Self-monitor for symptoms Avoid contact with vulnerable people Wear a mask during all social interactions Practice physical distancing of 2 meters where possible If symptoms develop see section 2	FOR HOUSEHOLD/SEXUAL CONTACTS OF A POSITIVE CASE: 5-day isolation from same date as the case Self-monitor for symptoms Days 6-10 continue to monitor for symptoms, avoid contact with vulnerable people, wear a mask during all social interactions, practice physical distancing of 2 meters where possible *it is not necessary to prolong isolation for each new positive case in the same family

^{1.} Severe immunosuppression: eg, solid organ transplant recipients, stem cell transplant recipients, hematological cancer undergoing chemotherapy)

^{*}Infection Prevention and Control.